



Sodium Hyaluronate



Hynadin

1 mg/mL (0.1% w/v)
Ophthalmic Solution (Eye Drops)
OCULAR LUBRICANT

FORMULATION:

Each mL contains:

Sodium hyaluronate 1 mg

PRODUCT DESCRIPTION:

Colorless and clear ophthalmic solution.

PHARMACEUTICAL DOSAGE FORM:

Ophthalmic solution

PHARMACODYNAMICS:

- Because sodium hyaluronate principle has the same structure as natural sodium hyaluronate and molecular weight of more than 4000 KDa, it contains excellent biocompatibility and viscoelasticity.
- Because sodium hyaluronate is produced through highly developed unique microbial fermentation process, it contains low amount of proteins and nucleic acids and is completely devoid of inflammatory substances.
- Because sodium hyaluronate contains high viscoelasticity, it shows good actions in maintaining cornea and protecting corneal endothelial cells.
- Sodium hyaluronate is a biological substance that has been shown in preclinical studies to accelerate the wound healing of corneal epithelium and to have water-holding properties. Its clinical usefulness has been established for the treatment of keratoconjunctival epithelial injury associated with dry eye syndrome or other disorders.

PHARMACOKINETICS:

Sodium hyaluronate are widely distributed in human tissue and intercellular fluid, including cataract fluid.

INDICATIONS:

For the treatment of keratoconjunctival injury associated with the following diseases: endogeneous diseases such as Sjogren's syndrome, Stevens-Johnson syndrome, dry eye syndrome, and exogeneous diseases caused by surgery, drugs, trauma, contact lens wearing and others.

DOSAGE AND ADMINISTRATION:

For ophthalmic use only.

Usually, instill 1 drop at a time to the eye 5-6 times daily. The dosage may be adjusted according to the patient's symptoms.

CONTRAINDICATIONS:

Patients known to be hypersensitive to any of the composition.

PRECAUTION:

At the time of administration, instruct the patient to be careful not to touch the tip of the bottle to the eye directly in order to avoid the contamination of the eye drops. This product must not be used while wearing soft contact lenses.

PREGNANCY AND LACTATION:

There are no adequate studies of using this drug in pregnant women and lactating mothers. This drug should be used during pregnancy or during lactating only if the potential benefit justifies the potential risk.

DRUG INTERACTIONS:

No report of drug interaction.

ADVERSE DRUG REACTIONS:

Hypersensitivity: Blepharitis, eyelid dermatitis.

Ophthalmic: Itching, eye irritation, corneal lesion, eg. keratitis superficial diffuse.

Inform your doctor in case of any adverse reactions related to drug use.

OVERDOSAGE:

No reports of overdosage.

EFFECTS ON ABILITY TO DRIVE AND OPERATE MACHINES:

The drug does not affect ability to drive and operate machines.

CAUTION:

Foods, Drugs, Devices, and Cosmetics Act prohibits dispensing without prescription.

"For suspected adverse drug reaction, report to the FDA: www.fda.gov.ph. Seek medical attention immediately at the first sign of any adverse drug reaction".

STORAGE CONDITIONS:

Store at temperatures not exceeding 30°C.

KEEP ALL MEDICINES OUT OF REACH OF CHILDREN.

USE TERM:

Three (3) years from manufacturing date.

Use drug within one (1) month from date of opening.

Do not use if the drug is out of date.

AVAILABILITY:

LDPE Plastic Bottle x 6 mL (Box of 1's)

DRP-7680-03

Date of First Authorization: November 24, 2021

Date of Revision of Package Insert: September 6, 2021

Manufactured by:

DAEWOO PHARM. CO., LTD.

153, Dadae-ro, Saha-gu, Busan,

Republic of Korea

Imported by:

BLISS PHARMA DISTRIBUTION

AND CONSULTANCY CORP.

A2, 236 San Jose Street, Barangay 2,

Carmona, Cavite

Distributed by:

MEDCHOICE VISION CARE INC.

Unit 907, 88 Corporate Center, Sedeño corner Valero St.,

Salcedo Village, Makati City, Metro Manila